

- Dunbar, R. M. 1988. *Primate Social Systems*. Croom Helm, Beckenham, UK.
- Fernandez-Duque, E., Corley, M. K. and Spence-Aizenberg, A. 2013. Family Aotidae (night monkeys). In: *Handbook of the mammals of the world, Volume 3: Primates*, R. A. Mittermeier, A. B. Rylands and D. E. Wilson (eds.), pp. 414–431. Lynx Edicions, Barcelona, Spain.
- Ferrari, S. F. and Beltrão-Mendes, R. 2011. Do snakes represent the principal threat to callitrichids? Fatal attack of a viper (*Bothrops leucurus*) on a common marmoset (*Callithrix jacchus*) in the Atlantic Forest of the Brazilian Northeast. *Primates* 52: 207–209.
- Garber, P. A. and Leigh S. R. 1997. Ontogenetic variation in small-bodied New World primates: Implications for patterns of reproduction and infant care. *Folia Primatol.* 68: 1–22.
- Goldizen, A.W. 1987. Tamarins and marmosets: communal care of offspring. In: *Primate Societies*, B. B. Smuts, D. L. Cheney, R. M. Seyfarth R. M., R. W. Wrangham and T. T. Struhsaker (eds.), pp. 34–43. University of Chicago Press, Chicago, USA.
- Heymann, E. W. 2000. The number of adult males in callitrichine groups and its implications for callitrichine social evolution. In: *Primate Males*, P. M. Kappeler (ed.), pp. 64–71. Cambridge University Press, Cambridge, UK.
- Huck, M., Lunenburg, M., Dávalos, V., Rotundo, M., DiFiore, A. and Fernandez-Duque, E. 2014. Double effort: parental behavior of wild Azara's owl monkeys in the face of twins. *Am. J. Primatol.* Early view.
- Knogge, C. and Heymann, E. W. 1995. Field observation of twinning in the dusky titi monkey, *Callicebus cupreus*. *Folia Primatol.* 65: 118–120.
- Lawrence, J. M. 2007. Understanding the pair bond in brown titi monkeys (*Callicebus brunneus*): male and female reproductive interests. Ph.D thesis, Columbia University, New York, USA.
- Link, A., Palma, A. C., Velez, A. and de Luna, A. G. 2006. Costs of twins in free-ranging white-bellied spider monkeys (*Ateles belzebuth belzebuth*) at Tinigua National Park, Colombia. *Primates* 47: 131–139.
- Ross, C. 1991. Life history pattern of New World monkeys. *Int. J. Primatol.* 12: 481–502.
- Rylands, A. B. and Mittermeier, R. A. 2013. Family Callitrichidae (marmosets and tamarins). In *Handbook of the mammals of the world, Volume 3: Primates*, R. A. Mittermeier, A. B. Rylands and D. E. Wilson (eds.), pp. 262–346. Lynx Edicions, Barcelona, Spain.
- Santana, M. M. 2012. Ecologia comportamental de um grupo de guigó-de-coimbra (*Callicebus coimbrai* Kobayashi and Langguth 1999) no leste de Sergipe. Masters dissertation, Universidade Federal de Sergipe, São Cristóvão, Brazil.
- Snowdon, C. T. 1996. Infant care in cooperatively breeding species. In: *Parental Care - Evolution, Mechanisms, and Adaptive Significance*, J. S. Rosenblatt and C. T. Snowdon (eds.), pp. 643–689. Academic Press, San Diego, USA.
- Souza-Alves, J. P. 2013. Ecology and Life-history of Coimbra-Filho's titi monkeys (*Callicebus coimbrai*) in the Brazilian Atlantic Forest. Doctoral thesis, Universidade Federal da Paraíba, João Pessoa. Brazil.
- Strier, K. B. 1991. Demography and conservation of an endangered primate, *Brachyteles arachnoides*. *Conserv. Biol.* 5: 214–218.
- Tardif, S. D. 1994. Relative energetic cost of infant care in small-bodied Neotropical primates and its relation to infant care patterns. *Am. J. Primatol.* 34: 133–143.
- Valeggia, C. R., Mendoza, S. P., Fernandez-Duque, E., Mason, W. A. and Lasley, B. 1999. Reproductive biology of female titi monkeys (*Callicebus moloch*) in captivity. *Am. J. Primatol.* 47: 183–195.
- Wright, P. C. 2013. *Callicebus* in Manu National Park: Territory, resources, scent marking and vocalizations. In: *Evolutionary Biology and Conservation of Titis, Sakis, and Uacaris*, L. M. Veiga, A. A. Barnett, S. F. Ferrari and M. A. Norconk (eds.), pp. 232–239. Cambridge University Press, Cambridge, UK.

---



---

## INFANTICIDES DURING PERIODS OF SOCIAL STABILITY: KINSHIP, RESUMPTION OF OVARIAN CYCLING, AND MATING ACCESS IN WHITE-FACED CAPUCHINS (*CEBUS CAPUCINUS*)

Valérie A. M. Schoof  
Eva C. Wikberg  
Katharine M. Jack  
Linda M. Fedigan  
Toni E. Ziegler  
Shoji Kawamura

### Introduction

In non-human primates, most infanticide cases (85%) occur during periods of social instability when a change-over of the top-ranking male occurs (van Schaik, 2000). In contrast, infanticides during periods of social stability have only been reported for a few species (Valderrama *et al.*, 1990; Murray *et al.*, 2007; Gibson *et al.*, 2008). The sexually selected infanticide (SSI) hypothesis (Hrdy 1974, 1979), proposes that infanticide is a male reproductive strategy in that males adopting this strategy benefit by gaining mating access to females who resume ovarian cycling prematurely following the death of their unweaned offspring. According to the SSI hypothesis, infanticide is a male reproductive strategy if: 1) the attacker is unrelated to the infant, 2) the mother's time to conception is shortened by the infant's death, and 3) the infanticidal male has an increased probability of siring the mother's future offspring. Though other hypotheses have been suggested to explain infanticide, most reported cases occur during or after periods of social instability and thus appear to fit the SSI hypothesis (Hrdy 1974, 1979; van Schaik, 2000).

In white-faced capuchins (*Cebus capucinus*), infanticides are commonplace during periods of social upheaval resulting

from alpha male replacements (Fedigan, 2003; Perry *et al.*, 2012; Jack *et al.*, 2014). Using a combination of behavioral, genetic, and hormone data, we examine whether the first confirmed infanticides during a socially stable period without immigrations or changes in the male dominance hierarchy fit the predictions of the SSI hypothesis.

## Methods

The study took place from July 2008 to November 2009 in the Santa Rosa Sector of the Area de Conservación Guanacaste, in northwestern Costa Rica (see Fedigan and Jack, 2012 for full description of the field site). Both confirmed infanticides occurred during observation of GN group by VAMS. Adult and subadult group composition was stable during the entire study period (4 adult males, 4 subadult males, 10 adult females, 12 juveniles, and 3-4 infants). We recorded data on male dominance interactions and socio-sexual interactions using *ad libitum* and 10-minute focal animal follows on all eight resident males (>623 hours of focal data; 2575 contact hours). *Ad libitum* behavioral data were collected at the time of each infanticide. While alpha male white-faced capuchins are easily recognizable, dominance relationships among subordinate males are difficult to discern because agonistic behaviors are rare (Schoof *et al.*, 2014). Therefore, we used the Elo-rating method to complement qualitative assessments of dominance rank, including that of the infanticidal male (Neumann *et al.*, 2011).

To identify maternal resumption of ovarian cycling from progesterone (P) and estradiol levels, we collected fecal samples from LV and RM once every three days, beginning immediately after the infanticide until each female was visibly pregnant (LV: Mar 17, 2009) or she disappeared/died (RM: May 19, 2009; see Carnegie *et al.*, 2011 for detailed methods). We defined the periovulatory phase as the day of the fP (fecal Progesterone) rise  $\pm$  3 days (Schoof *et al.*, 2014).

Using DNA extracted from infant tissue samples and adult fecal samples, we amplified nuclear DNA at 20 loci via PCR (Apm01, Ceb01, Ceb02, Ceb03, Ceb04, Ceb07, Ceb08, Ceb09, Ceb10, Ceb11, Ceb105, Ceb115, Ceb119, Ceb120, Ceb127, Ceb128, Ceb130, d3s1210, d7s794, pepl4; see Wikberg *et al.*, 2014 for detailed methods). Infant sires were assigned in CERVUS (Marshall *et al.*, 1998; Kalinowski *et al.*, 2007) at the 0.95 confidence level, and we used ML-RELATE (Kalinowski *et al.*, 2006) to calculate estimated relatedness values (*R*) and evaluate kin relationships among adults.

## Results

Case 1 - HW: At 17:55 on 30 August 2008, when the monkeys normally settle down, VAMS noticed something fall to the ground where some females were vocalizing nearby. After approaching slowly, an infant was observed



**Figure 1.** Photograph of lethal wound observed on HW (photo by Valerie A.M. Schoof). This characteristic wound was also observed on RM08's corpse.

(HW, 69 days old) lying on the ground, still breathing. VAMS immediately moved away so as not to interfere with the monkeys' behavior. The wounding incident was not directly observed, but it was assumed that a resident male was responsible because no extra-group individuals were in the area. On two occasions, the infant's mother (LV) came to the ground and made unsuccessful attempts to carry HW. Beginning at 18:15, in the dark, a single unidentified individual emitted lost calls (long distance contact calls) on and off for 30 min and moved away slightly for approximately 10 min before approaching again and being joined by a second monkey that emitted soft contact calls. At 18:19, the infant was no longer breathing. Examination of the remains confirmed the presence of a gash (~4 cm x 1 cm) in the lumbar/sacral region of the spinal cord, which was completely severed (Fig. 1). There were no other external wounds and no bones were visibly broken. Tissue samples were collected, and infant sex was confirmed as female.

Case 2 - RM08: During the afternoon of 24 January 2009, only the group's beta male (MM), a subordinate adult male, and two subadult subordinate males were present in the group. At 16:41, screams drew the attention of VAMS towards beta male MM and female RM, who were grappling, tangled and dangling by their tails. The beta male had RM's infant (RM08, 88 days old) in his mouth and seconds later he and RM08 dropped to the ground. The attack lasted no more than 20 seconds. The beta male was chased by RM and several other monkeys, and he disappeared out of sight. RM08 presumably died quickly thereafter because she did not make any further sounds or movements.

Approximately 15 minutes after the infanticide, beta male MM was again observed with the group and rested in proximity to a female and juvenile, and later carried an unidentified infant (>10 months of age). Subordinate adult male AD then approached the beta male and repeatedly engaged him in an aggressive coalitionary display against RM. The two males engaged in mutual play, AD sucked MM's tail, and then mounted him while duck facing and vocalizing. A few minutes later, the beta male was briefly chased by

RM and several others, including male AD, although no contact aggression was observed. Two additional mounts between MM and AD occurred at 17:45, with the males taking turns. RM stayed with the dead infant long after the rest of the group moved off to a sleep tree. She tried several times to carry the corpse up into the tree with her and was eventually successful, but left the infant perched on a tree limb before heading off in the group's direction. Examination of the remains confirmed the presence of a large gash (~7 cm × 2.5 cm) at the thoracic/lumbar intersection of the spinal cord (directly below the last rib), which was completely severed. There were no other external wounds and no bones were visibly broken. Tissue samples were collected, and infant sex was confirmed as female.

### Kinship

Alpha male BG was assigned as the sire to both HW and RM08. Based on demographic data and genetic analyses, alpha male BG and beta male MM were likely full-siblings ( $R=0.64$ ; full-siblings versus non-kin:  $P<0.001$ ; full-siblings versus half-siblings:  $P=0.01$ ; full-sibling versus parent-offspring  $P<0.001$ ). The beta male had relatively low  $R$ -values with both mothers (LV, HW's mother: 0.13; RM, RM08's mother: 0.03), but we could not distinguish if they were non-kin or half-siblings ( $P=0.19$  and  $P=0.11$  respectively). LV subsequently conceived and gave birth to an infant (HP) who was also sired by the alpha, rather than the beta male. Based on the estimated kinship between the beta male and the infants' parents, he was most likely the paternal uncle of the three infants.

### Sociosexual behavior and resumption of ovarian cycling

The conceptions of both HW and RM08 occurred prior to the start of this study, likely in January and May 2008 respectively (based on a mean gestation length of 157 days; Carnegie et al., 2011), and therefore we cannot evaluate male sexual access to the females at the time of conception. Prior to HW's infanticide, LV was observed copulating only with the alpha male BG. She had two periovulatory periods (POPs) beginning 88 and 109 days after the infanticide of HW. The second POP resulted in pregnancy and successful parturition of a subsequent infant on 22 May 2009, with an interbirth interval of 0.73 years (265 days). On 20 December 2008, during LV's conception POP, she and the alpha male engaged in sociosexual behavior (BG directed a duck face and dance display, behaviors that often lead up to a mount) and both engaged in sexual vocalizations but no copulations were observed. No other males were observed engaging in sociosexual behaviors with LV during this time. During the first two months of gestation, however, four other resident males engaged in sociosexual behaviors with LV, including beta male MM.

During RM08's gestation period, RM copulated (intromission observed) with a subordinate male and directed sexual vocalizations to the alpha male and was briefly mounted by him. In the four months following the infanticide and prior to her disappearance, sociosexual interactions – including

some copulations – were observed between RM and all resident males *except* infanticidal male MM. RM had two probable POPs beginning 49 and 83 days after RM08's infanticide, and her hormone profile is consistent with the progesterone and estradiol fluctuations observed in ovulating females. RM did not conceive again before her disappearance in late May 2009 (she was presumed dead at 26 years of age).

### Discussion

Predictions generated from the SSI hypothesis were only partially supported by the infanticides reported here. According to the SSI hypothesis, the attacker should be unrelated to the infant. Genetic data confirm that both infanticide victims were sired by the alpha male BG, rather than by beta male MM who was the known attacker of infant RM08. However, the alpha and beta male are likely full-siblings who transferred between groups together at least twice (Wikberg et al., 2014). To our knowledge no studies have reported paternal kin bias between males and their sibling's offspring, and males may occasionally kill related infants (Gibson et al., 2008; Clarke et al., 1994) if they have limited means to assess their relatedness to potential victims (Murray et al., 2007; Widdig, 2007). Instead, males may rely on their mating history with the mothers to assess kinship (Sugiyama, 1965; Hrdy, 1974; van Schaik, 2000). However, the killing of close kin's offspring suggests that if cognitive links between mating history and paternity probability exist, these are not transferable to the mating behavior of others.

The SSI prediction that the mother's time to conception would be shortened by the infant's death was generally supported. Capuchin infants are not fully weaned until 12 months (Carnegie et al., 2011; Fedigan and Jack, 2012). Given that both infants were under three months of age, the mothers were likely experiencing lactational amenorrhea at the time of the infanticides. Both mothers resumed ovarian cycling soon after the infanticide, and LV's interbirth interval was a mere 0.73 years, compared to the mean interval of 2.25 years for females whose infants survived to one year (Fedigan and Jack, 2012). The SSI hypothesis also predicts that the infanticidal male has an increased probability of siring the mother's future offspring. Following RM08's death, RM was observed engaging in sociosexual behavior with all resident males except the attacker MM. In the case of HW, the attacker was not identified, but the mother LV was observed copulating only with the alpha male during her conception period, and he sired the resulting offspring. Therefore, unless BG attacked his own infant (HW), the infanticidal male did not sire LV's subsequent infant. The probability of siring future offspring can also increase via an imminent or temporary increase in dominance rank, as was proposed for an observed infanticide in wedge-capped capuchins (Valderrama et al., 1990). We cannot discount this possibility since at the time of RM08's infanticide, the alpha male had been away for several hours.

The beta male may have committed infanticide despite a limited chance of siring the subsequent infant because the alpha male's absence may have minimized the risk of major repercussions. Alpha males and potential sires likely play an important role in infant defense as many infanticides occur after resident males, especially sires, disappear or die (van Schaik, 2000). While the role of protector male(s) is generally proposed as a response to threats from predators and non-resident conspecifics (van Schaik, 2000), potential sires may nonetheless protect infants from threats posed by co-resident males (Borries et al., 1999, Broom et al., 2004). Capuchin alpha males engage in infant defense during takeover attempts and intergroup encounters (Fedigan and Jack, 2013), and our analysis of long-term data shows that infanticide is just as likely to occur following rank reversals (where formerly subordinate males become alpha) as they are following takeovers by extragroup males (Jack et al., 2014). Although male vigilance is primarily directed towards predators and extragroup conspecifics (Rose and Fedigan, 1995), social vigilance appears to be related to the number of same-sex competitors (Jack, 2001). Thus, it is possible that alpha male absence, even if temporary, increases infanticide risk even in socially stable groups.

## Conclusions

Few studies have included the genetic and hormone data necessary to evaluate if the SSI hypothesis can explain infanticides by male primates during periods of social instability (e.g. Borries et al., 1999; Harris and Monfort 2003; Murray et al., 2007). The current study includes unique observations of infanticides combined with genetic and hormonal data, allowing us to evaluate whether the SSI hypothesis might also explain infanticides during a period of apparent social stability. The hormonal data supported the rapid resumption of ovarian cycling and a shortened interbirth interval as predicted by the SSI hypothesis. In contrast, behavioral data did not support increased reproductive access to females. Finally, the genetic data provided mixed support for the predictions of the SSI hypothesis; although the attacker did not sire the infant(s), he was likely closely related to their sire, and he did not sire the mother's subsequent infant. Taken together, the SSI hypothesis was not well supported by these infanticides during a socially stable period.

## Acknowledgements

We thank the administrators of the Área de Conservación Guanacaste (especially R. Blanco Segura); the graduate students and field assistants who helped with data collection (including E. Murillo, A. D. Melin, and N. Parr); F. Campos, A. Sato, M. Bergstrom, and T. Hiwatashi for help with genetic analyses; D. Wittwer for guidance with the hormone analyses; and C. A. Chapman and anonymous reviewers for helpful comments on this manuscript. VAMS was supported by FQRNT, Leakey Foundation, National Geographic Society, National Science Foundation, and

Tulane University's Stone Center for Latin American Studies (VAMS); additional funding received from Natural Sciences and Engineering Research Council of Canada (LMF), Canada Research Chairs Program (LMF), The Japan Society for Promotion of Science (ECW, SK); NIH support to the WNPRC [RR000167].

**Valérie A. M. Schoof**, Department of Anthropology, McGill University, 855 Sherbrooke Street West, Montreal, Quebec, Canada, H3A 2T7, E-mail: <vschoof@gmail.com; valerie.schoof@mcgill.ca> and Department of Anthropology, Tulane University, **Eva C. Wikberg**, Department of Integrated Biosciences, University of Tokyo, **Katharine M. Jack**, Department of Anthropology, Tulane University, **Linda M. Fedigan**, Department of Anthropology, University of Calgary, **Toni E. Ziegler**, Wisconsin National Primate Research Center, University of Wisconsin and **Shoji Kawamura**, Department of Integrated Biosciences, University of Tokyo

## References

- Borries, C., Launhardt, K., Epplen, C., Epplen, J. T. and Winkler, P. 1999. Males as infant protectors in Hanuman langurs (*Presbytis entellus*) living in multimale groups - defense pattern, paternity and sexual behaviour. *Behav. Ecol. Sociobiol.* 46: 350–356.
- Broom, M., Borries, C. and Koenig, A. 2004. Infanticide and infant defence by males - modelling the conditions in primate multi-male groups. *J. Theoret. Bio.* 231: 261–270.
- Carnegie, S. D., Fedigan, L. M. and Ziegler, T. E. 2011. Social and environmental factors affecting fecal glucocorticoids in wild, female white-faced capuchins (*Cebus capucinus*). *Am. J. Primatol.* 73:861–869.
- Clarke, M. R., Zucker, E. L. and Glander, K. E. 1994. Group takeover by a natal male howling monkey (*Alouatta palliata*) and associated disappearance and injuries of immatures. *Primates* 35:435–442.
- Fedigan, L. M. 2003. Impact of male takeovers on infant deaths, births and conceptions in *Cebus capucinus* at Santa Rosa, Costa Rica. *Int. J. Primatol.* 24:723–741.
- Fedigan, L. M. and Jack, K. M. 2012. Tracking monkeys in Santa Rosa: lessons from a regenerating tropical dry forest. In: *Long-term field studies of primates*, P.M. Kappeler and D. Watts (eds.), pp. 165–184. Springer Press.
- Fedigan, L. M. and Jack, K. M. 2013. Sexual conflict in white-faced capuchins: it's not whether you win or lose. In: *Evolution's Empress: Darwinian Perspectives on Women*, Fisher, M. L., Garcia, J. R., Chang, R. S. (eds), pp. 281–303. New York: Oxford University Press.
- Gibson, K. N., Vick, L. G., Palma, A. C., Carrasco, F. M., Taub, D. and Ramos-Fernandez, G. 2008. Intra-community infanticide and forced copulation in spider monkeys: A multi-site comparison between Cocha Cashu, Peru and Punta Laguna, Mexico. *Am. J. Primatol.* 70:485–489.
- Harris, T. R. and Monfort, S. L. 2003. Behavioral and endocrine dynamics associated with infanticide in a black

- and white colobus monkey (*Colobus guereza*). *Am. J. Primatol.* 61:135–142.
- Hrdy, S. B. 1974. Male-male competition and infanticide among the langurs (*Presbytis entellus*) of Abu Rajasthan. *Folia Primatol.* 22:19–58.
- Hrdy, S. B. 1979. Infanticide among animals: a review, classification, and examination of the implications for the reproductive strategies of females. *Ethol. Sociobiol.* 1:13–40.
- Jack, K. M. 2001. Effect of male emigration on the vigilance behavior of coresident males in white-faced capuchins (*Cebus capucinus*). *Int. J. Primatol.* 22: 715–732.
- Jack, K. M., Brasington, L. F., Ritchotte, K., Wikberg, E. C., Kawamura, S., and Fedigan, L. M. 2014. Bide your time: Method of alpha male replacement, infant deaths, and time to conception in *Cebus capucinus*. *Am. J. Primatol.* 76 (suppl 1): 79.
- Kalinowski, S. T., Wagner, A. P. and Taper, M. L. 2006. ML-RELATE: a computer program for maximum likelihood estimation of relatedness and relationship. *Mol. Ecol. Notes* 6:576–579.
- Kalinowski, S. T., Taper, M. L. and Marshall, T. C. 2007. Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Mol. Ecol.* 16:1099–1106.
- Marshall, T. C., Slate, J., Kruuk, L. E. B. and Pemberton, J. M. 1998. Statistical confidence for likelihood-based paternity inference in natural populations. *Mol. Ecol.* 7:639–655.
- Murray, C. M., Wroblewski, E. and Pusey, A. E. 2007. New case of intragroup infanticide in the chimpanzees of Gombe National Park. *Int. J. Primatol.* 28:23–37.
- Perry, S. 1998. Male-male social relationships in wild white-faced capuchins (*Cebus capucinus*). *Behav.* 135:139–172.
- Perry, S., Godoy, I. and Lammers, W. 2012. The Lomas Barbudal Monday Project: Two decades of research on *Cebus capucinus*. In: *Long-term field studies of primates*, P.M. Kappeler, D. Watts (eds.), pp.141–163. Springer Press.
- Rose, L. M., and Fedigan, L. M. 1995. Vigilance in white-faced capuchins, *Cebus capucinus*, in Costa Rica. *Behav.* 49:63–70.
- Schoof, V. A. M., Jack, K. M. and Ziegler, T. E. 2014. Male response to female ovulation in white-faced capuchins (*Cebus capucinus*): variation in fecal testosterone, dihydrotestosterone, and glucocorticoids. *Int. J. Primatol.* 35:643–660.
- Sugiyama, Y. 1965. On the social change of Hanuman langurs (*Presbytis entellus*) in their natural condition. *Primates* 6:381–418.
- Valderrama, X., Srikosamatara, S. and Robinson, J. G. 1990. Infanticide in wedge-capped capuchin monkeys, *Cebus olivaceus*. *Folia Primatol.* 54:171–176.
- van Schaik, C. P. 2000. Infanticide by male primates: The sexual selection hypothesis revisited. In: *Infanticide by males and its implications* C.P. van Schaik and C.H. Janson (eds.), pp. 27–60). Cambridge, UK: Cambridge University Press.
- Widdig, A. 2007. Paternal kin discrimination: the evidence and likely mechanisms. *Biol. Rev.* 82:319–334.
- Wikberg, E. C., Jack, K. M., Campos, F. A., Fedigan, L. M., Sato, A., Bergstrom, M. L., Hiwatashi, T. and Kawamura, S. 2014. The effect of male parallel dispersal on the kin composition of groups in white-faced capuchins (*Cebus capucinus*). *Anim. Behav.* 96: 9–17.

---



---

## PREDATION OF BIRDS BY AN ENDANGERED PRIMATE SPECIES, *CALLICEBUS COIMBRAI*, IN THE BRAZILIAN ATLANTIC FOREST

Luana Vinhas  
João Pedro Souza-Alves

The world faces a significant environmental crisis, in which continuous natural environments are being reduced to disturbed fragments (Ladle and Whittaker, 2011). Some species can take advantage of this process, but the majority of biodiversity is threatened by human activities (IUCN 2012). Ecological plasticity is regarded as a characteristic that favors species' survival when habitat becomes degraded (McKinney, 1997). Most primates have generalist diets and show some behavioral plasticity (Garber, 1987); however, current knowledge is concentrated on some well-studied species, while there is a lack of information for other primates.

Titi monkeys (*Callicebus* spp.) are regarded as primarily frugivorous primates, which complement their diets with invertebrate prey and other plant parts, such as leaves, seeds and flowers (Bicca-Marques and Heymann, 2013; DeLuycker, 2012; Heymann and Nadjafzadeh, 2013). These primates tolerate disturbed habitat (Heiduck, 2002; Jerusalinsky et al., 2006; Souza-Alves et al. 2011a) and it has already been suggested that titi monkeys might show some dietary plasticity. For example, Santos et al. (2012) report that *C. nigrifrons* can take advantage of temporarily available items, such as masting bamboos. Neri (1997) describes a male *C. personatus* driving a dove away from its nest and allowing the female to eat its egg. Souza-Alves et al. (2011) verified a high consumption of insects (i.e., caterpillar) during the dry season by *C. coimbrai*. However, up to now, there are no reports of titis preying on vertebrates.

Here, the predation of birds by one subadult *C. coimbrai* in a large fragment of Atlantic forest in the northeastern Brazil is reported. The observation appears to be the first record of predation of birds by *C. coimbrai* and by titis in general. The events were recorded at the largest fragment of the Mata do Junco Wildlife Refuge - MJWR (10°32'S, 37°03'W), which encompasses 522 ha of Atlantic Forest in the Northeastern Brazilian state of Sergipe, municipality of Capela, Brazil. Systematic monitoring of the *C. coimbrai* study group has occurred since 2011 until the present time (Chagas et al., 2013). Between January and March 2014

## Neotropical Primates

A Journal and Newsletter of the IUCN/SSC Primate Specialist Group

Vol. 21(2), December 2014

## Contents

### Articles

- Jealous of Mom? Interactions Between Infants and Adult Males During the Mating Season in Wild Squirrel Monkeys (*Saimiri collinsi*)  
*Luana V. P. Ruivo and Anita I. Stone*..... 165
- Reconocimiento de Primates y Amenazas para su Supervivencia en Bosques Pre- Montano y Montano de la Región Cajamarca, Perú  
*Rolando Aquino, Elvis Charpentier, Gabriel García, Iris Arévalo y Luis López*..... 171
- Census of the Blond Titi Monkey *Callicebus barbarabrownae* (Pitheciidae) in the Semi-Deciduous Atlantic Forest of Chapada Diamantina, Brazil  
*Cintia F. Corsini and Antonio Christian de A. Moura* ..... 177
- Emergence of Sex-Segregated Behavior and Association Patterns in Juvenile Spider Monkeys  
*Michelle A. Rodrigues*..... 183

### Short Articles

- Twinning in Titis (*Callicebus coimbrai*): Stretching the Limits of Biparental Infant Caregiving?  
*Marina M. de Santana, João Pedro Souza-Alves and Stephen F. Ferrari*..... 190
- Infanticides During Periods of Social Stability: Kinship, Resumption of Ovarian Cycling, and Mating Access in White-Faced Capuchins (*Cebus capucinus*) ..... 192  
*Valérie A. M. Schoof, Eva C. Wikberg, Katharine M. Jack, Linda M. Fedigan, Toni E. Ziegler and Shoji Kawamura*
- Predation of Birds by an Endangered Primate Species, *Callicebus coimbrai*, in the Brazilian Atlantic Forest..... 196  
*Luana Vinhas and João Pedro Souza-Alves*
- Observation of Weapon Use in a Group of Semi-Free Tufted Capuchins (*Sapajus* spp) ..... 199  
*Claire Hamilton, and Dorothy M. Fragaszy*
- Two Case Studies Using Playbacks to Census Neotropical Primates: *Callicebus discolor* and *Alouatta palliata aequatorialis* ..... 201  
*Andrea Salcedo R, Manuel Mejia, Katie Slocombe and Sarah Papworth*
- Report of a Black Spider Monkey (*Ateles chamek*) Swimming in a Large River in Central-Western Brazil..... 205  
*André Valle Nunes*
- Yellow-Tailed Woolly Monkey (*Oreonax flavicauda*: Humboldt 1812) Altitudinal Range Extension, Uchiza, Perú ..... 207  
*Néstor Allgas, Sam Shane, Ana Peralta and Noga Shane*
- Primatas da Reserva Particular do Patrimônio Natural Água Boa, Cacoal, Rondônia, Brasil..... 208  
*Almério Câmara Gusmão, Marcella Alves Crispim, Marcelo Lucian Ferronato e José de Sousa e Silva Júnior*
- Registro de *Callicebus cinerascens* (Spix, 1823) no médio Vale do Guaporé, Rondônia, Brasil..... 211  
*Almério Câmara Gusmão e Thatiane Martins da Costa*
- Predação ou Necrofagia de *Alouatta guariba clamitans* por *Leopardus pardalis*? ..... 212  
*William Douglas de Carvalho, Ayesha Ribeiro Pedrozo, Theany Biavatti, Luciana de Moraes Costa e Carlos Eduardo Lustosa Esbérard*
- Ocorrência de *Sapajus flavius* e *Alouatta belzebul* No Centro de Endemismo Pernambuco..... 214  
*Marcos de Souza Fialho, Mônica Mafra Valença-Montenegro, Thiago César Farias da Silva, Juliana Gonçalves Ferreira e Plautino de Oliveira Laroque*
- Capture of a Common Marmoset (*Callithrix jacchus*) by a Capuchin Monkey (*Sapajus* SP.) in the Ibura National Forest, Sergipe (Brazil)..... 219  
*Natasha M. Albuquerque and Saulo M. Silvestre, Thayane S. Cardoso, Juan Manuel Ruiz-Esparza, Patrício A. Rocha, Raone Beltrão-Mendes e Stephen F. Ferrari*

Recent Publications..... 221

Meetings ..... 226